



TITLE:

On ℓ^3 -divisibility of class numbers of ℓ -cyclic extensions (Algebraic Number Theory)

AUTHOR(S):

NAITO, Hirotada

CITATION:

NAITO, Hirotada. On ℓ^3 -divisibility of class numbers of ℓ -cyclic extensions (Algebraic Number Theory). 数理解析研究所講究録 1987, 603: 87-92

ISSUE DATE:

1987-01

URL:

<http://hdl.handle.net/2433/99653>

RIGHT:

On ℓ^3 -divisibility of class numbers of
 ℓ -cyclic extensions

by Hirotada NAITO

内藤 浩忠

The Faculty of Education

Kagawa University

Takamatsu, JAPAN

§ 1. Introduction.

In this talk, we consider a problem of the divisibility of class numbers of algebraic number fields of finite degree. Many people have studied the following problem:

Are there infinitely many algebraic number fields k satisfying some prescribed conditions whose class numbers are divisible by a given integer n ?

Kuroda[7] studied the case that fields k are imaginary quadratic fields in which a finite number of prescribed primes are ramified.

Yamamoto[11] studied the case that fields k are real quadratic fields.

Uchida[10] studied the case that fields k are cyclic extensions over the rational number fields \mathbb{Q} of degree 3.

Azuhata and Ichimura[1] studied the case that fields k have r_1 real places and r_2 imaginary places ($r_2 \geq 1$).

Nakano[9] generalized the above result to the case

including $r_2=0$.

Further references are found in Diaz y Diaz[3].

For a special class of quadratic number fields we know the following theorem.

THEOREM (Kaplan[6] and Yamamoto[12]) Let $p \equiv 3 \pmod{4}$ be the fixed prime. Let $q \equiv 1 \pmod{4}$ be the prime. Then the following properties are equivalent:

- (i) The class number of $\mathbb{Q}(\sqrt{-pq})$ is divisible by 8.
- (ii) The prime q is completely decomposed in K_8/\mathbb{Q} , where $K_8 = \mathbb{Q}(\sqrt{-1}, \sqrt[4]{p})$.

By Tchebotarev's density theorem we see that the density of the set consisting of the primes q with the above property (ii) is $1/8$. Therefore we may say that the density of $\mathbb{Q}(\sqrt{-pq})$ whose class numbers are divisible by 8 is $1/8$ for the fixed prime p . Cohn[2] called such a field K_8 the governing field and studied some types of quadratic fields.

In this talk, we investigate the ℓ^3 -divisibility of the cyclic extensions of degree ℓ , where ℓ is an odd prime number. Let p be the fixed prime such that $p \equiv 1 \pmod{\ell}$ or $p = \ell$. Let $q \equiv 1 \pmod{\ell}$ be a prime. We denote by L_p the cyclic extension over the rational number field \mathbb{Q} of degree ℓ where only the prime p is ramified. We denote by L_i ($1 \leq i \leq \ell-1$) the cyclic extensions over \mathbb{Q} of degree ℓ where only both of the primes p and q are

ramified. The class number of L_p is prime to ℓ and the index of the group of circular units of L_p in the unit group E of L_p is also prime to ℓ . Let ξ_1 and ξ_2 be circular units of L_p such that the images of the subgroups $\langle \xi_1 \rangle$ and $\langle \xi_1, \xi_2 \rangle$ in E/E^ℓ are invariant under the action of the Galois group of L_p/Q . We put $\mathcal{L}_p^2 = L_p(\zeta_\ell, \sqrt[\ell]{\xi_1})$ and $\mathcal{L}_p^3 = L_p(\zeta_\ell, \sqrt[\ell]{\xi_1}, \sqrt[\ell]{\xi_2})$, where ζ_ℓ is a primitive ℓ -th root of unity. Then we get:

THEOREM. For $r=2$ or 3 , the following properties are equivalent:

- (i) The class number of L_i is divisible by ℓ^r for some $1 \leq i \leq \ell-1$.
- (ii) The class number of L_i is divisible by ℓ^r for any $1 \leq i \leq \ell-1$.
- (iii) The prime q is completely decomposed in \mathcal{L}_p^r/Q .

REMARK 1. For $r=2$, this is a result of Inaba[4], c.f. also Gras[5]. In these papers it is shown that the property

(i) is equivalent to the property $\left(\frac{q}{p}\right)_\ell = \left(\frac{p}{q}\right)_\ell = 1$, where $\left(\frac{*}{*}\right)_\ell$ is the ℓ -th power residue symbol. We get $\mathcal{L}_p^2 = L_p(\zeta_\ell, \sqrt[\ell]{p})$ because of $\xi_1 = p\alpha^\ell$ for some $\alpha \in L_p$. Thus we see that (i) and (ii) are equivalent.

REMARK 2. If the class number of L_i is divisible by 3^3 , the ideal class group of L_i has an element of order 3^2 for $\ell=3$.

§ 2. The Proof.

We denote by $(\mathbb{Z}/m\mathbb{Z})^{\oplus s}$ the direct sum of s cyclic groups of order m . We see by class field theory that the property (iii) in Theorem 4 is equivalent to the following

(iii)' L_p has a $(\mathbb{Z}/\ell\mathbb{Z})^{\oplus r}$ -extension with conductor q .

At first we explain the case of $r=2$. Let H_c be the unramified $(\mathbb{Z}/\ell\mathbb{Z})^{\oplus 2}$ -extension over L_1 such that H_c/Q is a Galois extension. Then H_c/L_j ($j \neq 1$) is an unramified $(\mathbb{Z}/\ell\mathbb{Z})^{\oplus 2}$ -extension. Moreover H_c/L_p is a $(\mathbb{Z}/\ell\mathbb{Z})^{\oplus 2}$ -extension with conductor q .

Next we explain the case of $r=3$. We see by Proposition VI.6. in Gras[4] that the properties (i) and (ii) are equivalent.

We assume (ii). Let H_1/L_1 be the unramified extension whose Galois group is isomorphic to $(\mathbb{Z}/\ell\mathbb{Z})^{\oplus 3}$ for $\ell \geq 5$ or to $(\mathbb{Z}/3^2\mathbb{Z}) \oplus (\mathbb{Z}/3\mathbb{Z})$ for $\ell=3$ such that H_1/Q is a Galois extension. Let H be the compositum of H_i ($1 \leq i \leq \ell-1$). We see by the computation of the Galois group H/L_p that H contains the $(\mathbb{Z}/\ell\mathbb{Z})^{\oplus 3}$ -extension H_p/L_p with conductor q . Thus we get (iii)'.

We assume (iii)'. Let H_p/L_p be the $(\mathbb{Z}/\ell\mathbb{Z})^{\oplus 3}$ -extension with conductor q . Let H_c/L_p be the $(\mathbb{Z}/\ell\mathbb{Z})^{\oplus 2}$ -extension in H_p such that H_c is a Galois extension over Q . Let \mathfrak{P} be the prime ideal of L_p lying

over p . The prime ideal \mathfrak{P} is completely decomposed in H_c/L_p , because \mathfrak{P} is invariant under the action of the Galois group of L_p/Q . Let \mathfrak{P}_1 be the prime ideal of L_1 lying over p . We see that \mathfrak{P}_1 is completely decomposed in H_c/L_1 . We see that H_c/L_1 is an unramified $(\mathbb{Z}/\ell\mathbb{Z})^{\oplus 2}$ -extension. As \mathfrak{P}_1 is an element of genus group of L_1 , we see by class field theory that L_1 has an unramified abelian extension of degree ℓ^3 . Thus we get (i).

REFERENCES.

- [1] T. Azuhata and H. Ichimura : On the divisibility problem of the class numbers of algebraic number fields, J. Fac. Sci. Univ. Tokyo 30 (1984), 579-585.
- [2] H. Cohn : Introduction to the construction of class fields, Cambridge Univ. Press, 1985.
- [3] F. Diaz y Diaz : Sur le 3-rang des corps quadratique, Publ. Math. d'Orsay, N°78-11 (1978).
- [4] G. Gras : Sur les ℓ -classes d'idéaux dans les extensions cycliques relatives de degre premier ℓ , Ann. Inst. Fourier, Grenoble 23.3(1973),1-48, 23.4(1973),1-44.
- [5] E. Inaba : Uber die Struktur der ℓ -Klassengruppe zyklischer Zahlkorper vom Primzahlgrad ℓ , J. Fac.Sci. Univ. Tokyo 4 (1940), 61-115.

- [6] P. Kaplan : Divisibilité par 8 du nombre des classes des corps quadratiques dont le 2-groupe des classes est cycliques, et réciproque biquadratique, J. Math. Soc. Japan 25 (1973), 596-608.
- [7] S.-N. Kuroda : On the class number of imaginary quadratic number fields, Proc. Japan Acad. 40 (1964), 365-367.
- [8] H. Naito : Some results on class numbers and unramified extensions of algebraic number fields, the 19th Taniguchi Symposium (1986).
- [9] S. Nakano : On ideal class group of algebraic number fields, J. Reine Angew. Math. 358 (1985), 61-75.
- [10] K. Uchida : Class numbers of cubic cyclic fields, J. Math. Soc. Japan 26 (1974), 447-453.
- [11] Y. Yamamoto : On unramified Galois extensions of quadratic number fields, Osaka J. Math. 7 (1970), 57-76.
- [12] Y. Yamamoto : Divisibility by 16 of class number of quadratic fields whose 2-class groups are cyclic, Osaka J. Math. 21 (1984), 1-22.